

VI. POLLUTANT LOADINGS

A. INTRODUCTION

Pollutant loadings from point and nonpoint sources of pollution to the St. Louis River AOC are discussed in this chapter. Most of the information used to calculate pollutant loadings was taken from discharge monitoring reports for point source dischargers in Minnesota and Wisconsin. No estimate of nonpoint loadings were done at this time due to the limited data available and the variety of sources. Nonpoint sources of pollution include runoff from urban areas, construction sites, industrial sites, and agricultural land. The point sources of pollution can be traced to a pipe or outfall from a municipal or industrial facility.

1. Point Sources

There are four major discharges to the St. Louis River AOC of which two are municipal sewage treatment plants and two are industrial facilities. The discharges from these facilities are described in Table VI.1. Loadings are broken down by Biochemical Oxygen Demand (BOD), Total Suspended Solids (TSS), and Phosphorous. Average monthly discharge flows are given in million gallons per day (mgd). The tables are based on discharge monitoring reports from 1989 and 1990. These reports list the average monthly discharge loadings and flows that are monitored periodically by each facility.

Total pollutant loadings for conventional pollutants from all point sources are described in Table VI.2. This includes all permitted facilities that monitor for conventional pollutants. Loadings were calculated from monthly averages so the actual loadings could vary significantly in a given period from estimates given in the report. Four of these facilities do not have a continuous discharge throughout the year so the loadings listed are estimated based on the months they actually discharged. Facilities that discharge noncontact cooling water only have not been included in this report.

Pollutant loadings for toxic parameters have been estimated for those parameters where there are sufficient data (Table VI.3). Loadings estimates from point sources were not made for parameters with only one or two samples. Priority pollutant scans for permit reissuance were a major source of these data. In many cases however, there have been only one or two recent tests for a toxic parameter. Past discharge monitoring for heavy metals and most organics has not been done on a continuous basis, consequently an accurate estimate cannot be made at this time for many parameters. The process of permit reissuance is underway for Murphy Oil USA, The City of Superior, and Superior Fiber Products, the three major dischargers on the Wisconsin side of the AOC. These permits will require monitoring for and will limit discharge of many parameters, including bioaccumulating substances.

Mercury is known to be a concern in the discharge from the Western Lake Superior Sanitary District. Elevated concentrations have been found in the water and sediments of the St. Louis River near the discharge from the WLSSD. This discharge is due to mercury removal from the incinerator stack by a wet scrubbing process. The discharge permit for WLSSD was reissued in August 1990 and will require continuous monitoring for a range of toxic parameters.

The WLSSD Regional plant in Duluth was constructed in the late 1970's to treat waste from virtually all the industrial and municipal discharges in the area. Since that time dramatic improvements have occurred in general water quality as a result of BOD and Phosphorus reductions to the river. Concentrations of toxics in the sediments and water column are still at levels of concern leading to fish consumption advisories and the biological effects.

2. Nonpoint Sources

The St. Louis River AOC receives nonpoint pollution from a variety of sources due to the many types of land uses in the watershed. The most obvious source is the red clay erosion from tributaries such as the Nemadji River, Pokegama, Little Pokegama, and Red Rivers. These rivers contribute a seasonally heavy load of sediments to the St. Louis River and Superior Bay causing turbidity and sedimentation. The Red Clay project study conducted in the 1970's found that a large amount of erosion was due to naturally occurring bank slumpage. However, land use practices in the watersheds can contribute significantly to erosion. In addition to sediment input it was found that nutrients can attach to sediment particles causing excess nutrient loading.

The heavy concentration of industries located adjacent to the River and Bay on both the Wisconsin and Minnesota side contribute a variety of pollutants through storm water runoff. This type of runoff typically contains heavy metals, volatile organic chemicals (VOCs), oil, grease and a variety of organic and inorganic contaminants. Research has shown that shock loadings from stormwater events can have a greater effect than point source discharges. Storm sewers from Duluth and Superior discharge to the St. Louis River contributing the types of contaminants previously mentioned. Pollutant loadings from the Superior combined overflow #2 are described in Table VI.1. This indicates the significance of stormwater runoff in the area.

Table VI.1 Monthly Average Loadings for 1989-1990 from Facilities Discharging to the St. Louis River AOC

Western Lake Superior Sanitary District Effluent - 1989-90 - Average

| <u>Month</u> | <u>Flow (mgd)</u> | | <u>*BOD₅(lbs/day)</u> | | <u>TSS (lbs/day)</u> | | <u>Total Phosphorus (lbs/day)</u> | |
|--------------|-------------------|-------------|----------------------------------|-------------|----------------------|-------------|-----------------------------------|-------------|
| | <u>Ave.</u> | <u>Max.</u> | <u>Ave.</u> | <u>Max.</u> | <u>Ave.</u> | <u>Max.</u> | <u>Ave.</u> | <u>Max.</u> |
| January | 29.61 | 33.57 | 1053 | 1251 | 2475 | 3077 | 183 | 380 |
| February | 29.64 | 32.93 | 1155 | 1444 | 2100 | 2792 | 178 | 305 |
| March | 35.09 | 72.49 | 1406 | 2820 | 1906 | 3915 | 158 | 462 |
| April | 42.17 | 77.33 | 2011 | 4263 | 3789 | 8699 | 159 | 717 |
| May | 38.24 | 61.47 | 1490 | 2380 | 2141 | 5427 | 164 | 1343 |
| June | 36.12 | 42.15 | 1554 | 2116 | 2850 | 4669 | 193 | 366 |
| July | 32.54 | 40.61 | 1501 | 1974 | 4702 | 6954 | 201 | 551 |
| August | 36.82 | 76.68 | 1985 | 3524 | 6289 | 11827 | 262 | 889 |
| September | 42.37 | 94.27 | 1523 | 3299 | 6067 | 20,253 | 242 | 1480 |
| October | 38.00 | 55.16 | 1261 | 2759 | 1878 | 4254 | 203 | 390 |
| November | 32.99 | 37.03 | 996 | 1317 | 1514 | 2606 | 1397 | 341 |
| December | 30.76 | 33.12 | 1009 | 1239 | 1187 | 1665 | 190 | 282 |
| Ave. | 35.36 | 54.70 | 1412 | 2353 | 3075 | 6345 | 294 | 626 |

*Reported as Carbonaceous BOD

**Table VI.1 cont. Monthly Average Loadings for 1989-1990 from Facilities
Discharging to the St. Louis River AOC**

City of Superior Effluent - 1989-1990 - Average

| <u>Month</u> | <u>Flow (mgd)</u> | | <u>BOD₅(lbs/day)</u> | | <u>TSS (lbs/day)</u> | | <u>Total Phosphorus (lbs/day)</u> | |
|--------------|-------------------|-------------|---------------------------------|-------------|----------------------|-------------|-----------------------------------|-------------|
| | <u>Ave.</u> | <u>Max.</u> | <u>Ave.</u> | <u>Max.</u> | <u>Ave.</u> | <u>Max.</u> | <u>Ave.</u> | <u>Max.</u> |
| January | 3.49 | 4.20 | 426 | 1081 | 303 | 791 | 16 | 39 |
| February | 3.63 | 4.46 | 498 | 1066 | 256 | 647 | 15 | 42 |
| March | 3.96 | 5.43 | 629 | 1399 | 349 | 1205 | 16 | 38 |
| April | 4.03 | 5.48 | 525 | 1481 | 326 | 1193 | 11 | 40 |
| May | 4.12 | 5.02 | 658 | 1466 | 378 | 1621 | 13 | 44 |
| June | 3.58 | 5.10 | 669 | 1504 | 251 | 1594 | 9 | 40 |
| July | 3.61 | 4.88 | 695 | 1820 | 291 | 1253 | 12 | 37 |
| August | 3.74 | 4.99 | 478 | 1230 | 203 | 864 | 10 | 29 |
| September | 4.40 | 5.23 | 418 | 1001 | 272 | 618 | 12 | 32 |
| October | 3.77 | 4.83 | 388 | 1323 | 226 | 749 | 9 | 31 |
| November | 3.45 | 4.43 | 549 | 1394 | 202 | 684 | 9 | 22 |
| December | 2.51 | 3.21 | 440 | 930 | 134 | 301 | 9 | 26 |
| Ave. | 3.68 | 4.77 | 531 | 1308 | 266 | 960 | 12 | 35 |

**Table VI.1 cont. Monthly Average Loadings for 1989-1990 from Facilities
Discharging to the St. Louis River AOC**

City of Superior Effluent - Combined Sewer Overflow #2 - 1989-90 - Average

| <u>Month</u> | <u>Flow (mgd)</u> | | <u>BOD₅(lbs/day)</u> | | <u>TSS (lbs/day)</u> | | <u>Total Phosphorus (lbs/day)</u> | |
|--------------|-------------------|-------------|---------------------------------|-------------|----------------------|-------------|-----------------------------------|-------------|
| | <u>Ave.</u> | <u>Max.</u> | <u>Ave.</u> | <u>Max.</u> | <u>Ave.</u> | <u>Max.</u> | <u>Ave.</u> | <u>Max.</u> |
| January | 4.60 | | 544 | | 822 | | 22 | |
| February | ND | -- | -- | -- | -- | -- | -- | -- |
| March | 12.02 | 18.28 | 1865 | 4028 | 3488 | 9031 | 85 | 232 |
| April | 10.76 | 20.56 | 1107 | 3143 | 1800 | 4876 | 48 | 110 |
| May | 9.10 | 14.78 | 989 | 3314 | 1247 | 4736 | 29 | 63 |
| June | 7.67 | 9.49 | 748 | 1019 | 889 | 1379 | 13 | 20 |
| July | 8.28 | 10.60 | 424 | 906 | 1276 | 4723 | 15 | 21 |
| August | 19.08 | 32.80 | 3346 | 11,727 | 5952 | 15,628 | 85 | 216 |
| September | 11.33 | 42.34 | 904 | 3061 | 1972 | 9351 | 51 | 164 |
| October | 13.54 | 30.10 | 1119 | 1866 | 2998 | 5204 | 60 | 96 |
| November | 8.79 | -- | 440 | -- | 807 | -- | 45 | -- |
| December | ND | -- | -- | -- | -- | -- | -- | -- |
| Ave. | 10.52 | 18.47 | 1149 | 2961 | 2125 | 5575 | 44 | 94 |

ND - No Discharge

NOTE: The averages represented in this table are calculated only for the days of actual discharge. These discharges occur intermittently. The averages would be significantly lower if they were calculated as an average daily discharge on a monthly basis which assumes 30 days of discharge per month.

**Table VI.1 Cont. Monthly Average Loadings for 1989-1990 from Facilities
Discharging to the St. Louis River AOC**

Superior Fiber Products Effluent - 1989-1990 Average

| <u>Month</u> | <u>Flow (mgd)</u> | | <u>BOD₅ (lbs/day)</u> | | <u>TSS (lbs/day)</u> | |
|--------------|-------------------|-------------|----------------------------------|-------------|----------------------|-------------|
| | <u>Ave.</u> | <u>Max.</u> | <u>Ave.</u> | <u>Max.</u> | <u>Ave.</u> | <u>Max.</u> |
| January | .667 | .714 | 2633 | 4386 | 197 | 369 |
| February | .667 | .712 | 2836 | 3786 | 109 | 360 |
| March | .697 | .734 | 1880 | 3038 | 168 | 259 |
| April | .677 | .733 | 1980 | 5764 | 193 | 258 |
| May | .657 | .742 | 2161 | 5225 | 158 | 331 |
| June | .668 | .697 | 2760 | 4327 | 222 | 331 |
| July | .685 | .750 | 2545 | 4547 | 205 | 602 |
| August | .697 | .745 | 2320 | 4166 | 182 | 338 |
| September | .680 | .740 | 2441 | 5114 | 209 | 482 |
| October | .688 | .741 | 2531 | 3792 | 218 | 471 |
| November | .665 | .735 | 2755 | 3905 | 204 | 286 |
| December | .665 | .721 | 2843 | 5389 | 190 | 324 |
| Average | .676 | .730 | 2473 | 4453 | 188 | 368 |

**Table VI.1 Cont. Monthly Average Loadings for 1989-1990 from Facilities
Discharging to the St. Louis River AOC**

Murphy Oil Company Effluent - 1989-1990 Average

| <u>Month</u> | <u>Flow (mgd)</u> | | <u>BOD₅ (lbs/day)</u> | | <u>TSS (lbs/day)</u> | |
|--------------|-------------------|-------------|----------------------------------|-------------|----------------------|-------------|
| | <u>Ave.</u> | <u>Max.</u> | <u>Ave.</u> | <u>Max.</u> | <u>Ave.</u> | <u>Max.</u> |
| January | .302 | .424 | 72 | 106 | 25 | 42 |
| February | .209 | .252 | 64 | 82 | 16 | 29 |
| March | .322 | .596 | 77 | 128 | 30 | 61 |
| April | .376 | .896 | 76 | 163 | 29 | 72 |
| May | .340 | .721 | 61 | 167 | 24 | 57 |
| June | .319 | .576 | 43 | 90 | 16 | 37 |
| July | .280 | .520 | 35 | 84 | 18 | 57 |
| August | .314 | .864 | 36 | 61 | 19 | 55 |
| September | .388 | .900 | 31 | 74 | 31 | 148 |
| October | .313 | .632 | 93 | 105 | 23 | 60 |
| November | .221 | .480 | 69 | 83 | 19 | 30 |
| December | .237 | .328 | 86 | 97 | 22 | 42 |
| Average | .302 | .599 | 62 | 103 | 23 | 58 |

Table VI.2 Total Annual Loadings of Conventional Pollutants from Point Sources to the St. Louis River AOC for 1989 and 1990

Average Annual Loadings in lbs.

| <u>Facility</u> | <u>BOD₅</u> | <u>Total Suspended Solids</u> | <u>Total Phosphorus</u> | <u>Ave. Monthly Discharge (mgd) Flow</u> |
|---|------------------------|-------------------------------|-------------------------|--|
| Western Lake Superior Sanitary District | 515,380 | 1,122,375 | 107,310 | 35.36 |
| City of Superior | 193,724 | 96,937 | 4,121 | 3.68 |
| City of Superior CSO) #2 | 50,468 | 88,441 | 1,816 | 10.52 ⁽¹⁾ |
| Superior Fiber Products | 901,886 | 72,270 | -- | .676 |
| Murphy Oil | 22,630 | 8,212 | -- | .302 |
| Burlington Northern ⁽¹⁾ | -- | 953 | -- | .16 ⁽²⁾ |
| Chicago & Northwestern ⁽²⁾ | -- | 395 | -- | .019 ⁽³⁾ |
| Duluth Winnipeg and Pacific | 1017 | 153 | -- | .006 |
| Village of Superior ⁽³⁾ | 146 | 630 | -- | .279 ⁽⁴⁾ |

⁽¹⁾This represents the average flow when the facility is actually discharging. Discharge occurred approximately 37 days each year in 1989 and 1990.

⁽²⁾Burlington Northern discharged in 7 months each year in 1989-90. This represents an average flow for those months.

⁽³⁾Chicago and Northwestern discharged 8 months each year in 1989-90. This represents an average flow for those months.

⁽⁴⁾The Village of Superior operates a fill and draw lagoon system which is usually drained in the spring and fall. Since no discharge occurred in 1989 this represents the average flow of the two drawdowns in 1990 over a 16 day and 28 day period.

**Table VI.3 Annual Mass Loadings Estimates - Toxic Substances
Point Source Discharges Into the St. Louis River AOC**

| <u>Facility</u> | <u>Flow mgd</u> | <u>Dates of Samples</u> | <u>Substance</u> | <u>Estimated lbs/yr discharge</u> | <u>Total number samples</u> |
|-----------------|---------------------|-------------------------|--|-----------------------------------|-------------------------------|
| Murphy Oil | .285 ave. discharge | 2/89 - 9/90 | Cr ⁺³ | 2.8 | 174 |
| | | | Cr ⁺⁶ | <0.7 | 174 |
| | | | Cu | 5.5 | 25 |
| | | | Ni | 8.5 | 25 |
| | | | Zn | 30.1 | 25 |
| | | | CN | 97.2 | 25 |
| | | | total phenols | 43.8 | 104 (from 2C application) |
| | | | Hg | (0.06) ¹ | 25 (3 detects) |
| | | | PCBs | no data | |
| | | | dioxin* | ---- | *detected in process |
| | | | pentachlorophenol | ---- | 1 |
| | | | NH ₃ -N | 588 | 174 |
| | | | ¹ Hg loading estimate very tenuous, sporadic detections | | |
| Superior Fiber | .68 ave. discharge | 7/86 - 1988 | Hg | 1.28 | 19 (13 detects) |
| | | | Zn | 255. | 32 |
| | | | Cu | 21.7 | 30 (29 detects) |
| | | | PCBs | not detected | 1 |
| | | | dioxin | no data | |
| | | | pentachlorophenol | not detected | 1 |
| | | | total phenols | 994. | 12 (from 2C application-1986) |

**Table VI.3 cont. Annual Mass Loadings Estimates - Toxics Substances
Point Source Discharges Into the St. Louis River AOC**

| <u>Facility</u> | <u>Flow mgd</u> | <u>Dates of Samples</u> | <u>Substances</u> | <u>Estimated lbs/yr discharge</u> | <u>Total number samples</u> |
|---------------------|----------------------|-------------------------|---|-----------------------------------|-----------------------------|
| City of Superior | 5 mgd design flow | 1990 - 91 | Insufficient data to estimate loadings 1990 Priority pollutant scan: 1 set metals data, organics pending | | |
| Burlington Northern | .0645 ave. discharge | 4/88-10/89 | Cu | 0.29 | 17 (4 detects) |
| | | | Zn | 0.77 | 17 (9 detects) |
| | | | Cd | not detected | 3 |
| | | | Cr | not detected | 3 |
| | | | Pb | not detected | 3 |
| WLSSD | 36 mgd av. discharge | | Cd | 324.85 | |
| | | | Cr | 2555 | |
| | | | Cu | 3452.9 | |
| | | | Pb | 4164.65 | |
| | | | Hg | 21.9 | |
| | | | Ni | 3766.8 | |
| | | | Se | 3682.85 | |
| | | | Zn | 10227.30 | |
| | | | Acetone | 1825 | |
| | | | 2-Butanone | 1095 | |
| | | | Carbon Disulfide | 2190 | |
| | | | 2,4-Dimethylphenol | 0 | |
| | | | Phenol | 262.8 | |
| | | | Benzyl Alcohol | 365 | |
| | | | Benzoic Acid | 2190 | |
| | | | 2,4,5-Trichlorophenol | 365 | |
| | | | 2,4,6-Trichlorophenol | 0 | |
| | | | Bis(2-ethylhexyl) Phthalate | 7544.55 | |
| | | | Butylbenzyl Phthalate | 0 | |
| | | | Chloroform | 11300.4 | |

| | |
|--------------------|------------|
| Methylene Chloride | 1752 |
| Hepychlor Epoxide | 0 |
| 2,3,7,8 TCDD | 0.00006205 |
| 2,3,7,8 TCDF | 0.0000949 |

Figure VI.1 Biochemical Oxygen Demand Loadings from Major Point Sources

